

Supplementary Material

Supplementary Table 1. Baseline characteristics of male participants based on their smoking status.

Data are presented as mean \pm standard deviation or as a percentage (%).

	Overall	Non-smoker	Waterpipe	Cigarettes	Dual	P value
	n=551	n=293	n=46	n=161	n=51	
Height (m)	167.3 \pm 9.3	163.6 \pm 9.2	171.2 \pm 8.6	171.0 \pm 6.8	173.9 \pm 8.0	<0.001
Weight (kg)	74.7 \pm 17.2	66.3 \pm 12.0	86.5 \pm 18.6	83.1 \pm 17.2	86.3 \pm 16.0	<0.001
BMI (kg/m ²)	26.51 \pm 4.8	24.7 \pm 3.4	29.4 \pm 5.2	28.4 \pm 5.5	28.5 \pm 4.8	<0.001
BMI Categories						<0.001
(%)						
• Underweight	4.0%	6.1%	0%	1.9%	2.0%	
• Normal	32.8%	43.3%	15.2%	23.6%	17.6%	
• Overweight	45.9%	50.5%	47.8%	37.9%	43.1%	
• Obese	17.2%	0%	37.0%	36.6%	37.3%	
Lean mass (kg)	45.7 \pm 10.8	40.0 \pm 9.4	52.8 \pm 9.9	51.7 \pm 7.7	53.5 \pm 8.0	<0.001
Fat mass (kg)	26.3 \pm 9.5	23.8 \pm 6.6	30.7 \pm 11.7	28.4 \pm 11.3	29.7 \pm 10.5	<0.001
Body fat (%)	34.9% \pm 8.0	36.0% \pm 8.1	34.9% \pm 7.5	33.2% \pm 7.7	33.6% \pm 7.8	0.003

Supplementary Table 2. Baseline characteristics of female participants based on their smoking status.

Data are presented as mean \pm standard deviation or as a percentage (%).

	Overall	Non-smoker	Waterpipe	Cigarettes	Dual	P value
	n=328	n=205	n=62	n=35	n=26	
Height (m)	167.4 \pm 9.3	164.1 \pm 9.0	172.4 \pm 7.5	173.9 \pm 6.4	173.3 \pm 5.9	<0.001
Weight (kg)	74.4 \pm 18.5	65.3 \pm 12.5	91.1 \pm 17.4	88.9 \pm 15.6	86.5 \pm 17.2	<0.001
BMI (kg/m ²)	26.3 \pm 5.0	24.1 \pm 3.4	30.7 \pm 5.6	29.3 \pm 4.6	28.7 \pm 5.0	<0.001
BMI Categories						<0.001
(%)						
• Underweight	4.6%	7.3%	0%	0%	0%	
• Normal	36.3%	45.9%	17.7%	14.3%	34.6%	
• Overweight	42.4%	46.8%	32.3%	48.6%	23.1%	
• Obese	16.8%	0%	50.0%	37.1%	42.3%	
Lean mass (kg)	45.0 \pm 11.2	39.9 \pm 9.5	53.5 \pm 8.2	54.4 \pm 7.7	53.2 \pm 8.4	<0.001
Fat mass (kg)	26.6 \pm 10.2	22.9 \pm 6.6	34.5 \pm 12.5	31.3 \pm 11.1	30.2 \pm 12.2	<0.001
Body fat (%)	35.4% \pm 8.0	35.2% \pm 8.1	37.0% \pm 7.9	34.7% \pm 7.2	34.0% \pm 8.8	0.3