

Electronic cigarette use and its association with asthma, chronic obstructive pulmonary disease (COPD) and asthma-COPD overlap syndrome among never cigarette smokers

Emine Bircan¹, Ummugul Bezirhan², Austin Porter^{3,4}, Pebbles Fagan^{5,6}, Mohammed S. Orloff^{1,7}

ABSTRACT

INTRODUCTION Although smoking is a strong risk factor for lung diseases including asthma, COPD, and asthma-COPD overlap syndrome (ACOS), studies are needed to examine the association between e-cigarettes and asthma, COPD, and ACOS. This study evaluated the association between e-cigarette use and self-reported diagnosis of asthma, COPD, and ACOS using a large nationally representative sample of adults aged ≥ 18 years in the United States.

METHODS Cross-sectional data from the Behavioral Risk Factor Surveillance System (BRFSS) from 2016 to 2018 was used to examine self-reported information on current e-cigarette use, demographic variables, and asthma and COPD status among never cigarette smokers ($n=8736$). Asthma and COPD were measured by self-reported diagnosis, and respondents who reported having both diagnoses were then classified as having ACOS. Of the 46079 never cigarette smokers, 4368 non-e-cigarette smokers were 1:1 propensity score-matched to e-cigarette smokers on age, sex, race/ethnicity and education level. We used multinomial logistic regression to examine association between current e-cigarette use and self-report asthma, COPD, and ACOS while controlling for marital status and employment in addition to matching variables.

RESULTS Compared with never e-cigarette smokers, e-cigarette smokers had increased odds of self-reported ACOS (OR=2.27; 95% CI: 2.23–2.31), asthma (OR=1.26; 95% CI: 1.25–1.27) and COPD (OR=1.44; 95% CI: 1.42–1.46).

CONCLUSIONS Data from this large nationally representative sample suggest that e-cigarette use is associated with increased odds of self-reported asthma, COPD, and ACOS among never combustible cigarette smokers. The odds of ACOS were twice as high among e-cigarette users compared with never smokers of conventional cigarettes. The findings from this study suggest the need to further investigate the long-term and short-term health effects of e-cigarette use, since the age of those at risk in our study was 18–24 years.

AFFILIATION

- 1 Department of Epidemiology, Fay W. Boozman College of Public Health, University of Arkansas for Medical Sciences, Little Rock, United States
- 2 Teachers College, Columbia University, New York, United States
- 3 Department of Health Policy and Management, Fay W. Boozman College of Public Health, University of Arkansas for Medical Sciences, Little Rock, United States
- 4 Arkansas Department of Health, Little Rock, United States
- 5 Department of Health Behavior and Health Education, Fay W. Boozman College of Public Health, University of Arkansas for Medical Sciences, Little Rock, United States
- 6 Center for the Study of Tobacco, Department of Health Behavior and Health Education, Fay W. Boozman College of Public Health, University of Arkansas for Medical Sciences, Little Rock, United States
- 7 Winthrop P. Rockefeller Cancer Institute, University of Arkansas for Medical Sciences, Little Rock, United States

CORRESPONDENCE TO

Mohammed S. Orloff. Department of Epidemiology, Fay W. Boozman College of Public Health, University of Arkansas for Medical Sciences, 4301 West Markham Street, Slot #820, Little Rock, AR 72205-7199, United States. E-mail: MSOrloff@uams.edu

KEYWORDS

COPD, e-cigarettes, combustible cigarettes, asthma, asthma-COPD overlap syndrome (ACOS)

Received: 26 November 2020

Revised: 23 January 2021

Accepted: 25 January 2021

INTRODUCTION

Overall tobacco use has declined in the United States over the past 50 years, but the prevalence of

electronic cigarette (e-cigarette) use has increased rapidly in United States since their introduction in 2007¹. E-cigarettes are battery-powered devices that

create an aerosol by heating up liquid that usually includes nicotine, flavorings, propylene glycol, vegetable glycerin and other chemicals². The tobacco industry introduced e-cigarettes as a safer alternative to tobacco smoking³. Around 8.1 million (3.2%) US adults reported using e-cigarettes in 2018; the prevalence of e-cigarette use increased from 2.8% to 3.2% during 2017–2018⁴.

The causal relationship between cigarette smoking and respiratory diseases is well documented⁵. Over 80% of chronic obstructive pulmonary disease (COPD) cases are caused by cigarette smoking and cigarette smoking exacerbates asthma in adults⁵. The respiratory health effects of e-cigarette use are still not clear. Several studies that have examined the negative short-term health effects of e-cigarette use found that people who had smoked e-cigarettes for several months experienced an increase in shortness of breath and coughs^{2,6}. A recent consensus report from the National Academies of Science, Engineering, and Medicine on the Public Health Consequences of E-cigarettes⁷ showed moderate evidence that e-cigarettes increase coughing, wheezing as well as asthma exacerbation in adolescents. There is little evidence showing improvement in lung function and respiratory symptoms among adult smokers with asthma who switch to e-cigarettes completely or partly (dual use)⁷.

Data on the relationship between e-cigarette use and respiratory diseases such as COPD and asthma are scarce. Existing studies suggest that e-cigarette use is associated with chronic bronchitis, emphysema, COPD⁸ and asthma⁹. COPD and lifetime asthma are diagnosed in approximately 6% and 13% of the US adult population, respectively^{10,11}. Although it varies by population, prevalence of asthma-COPD overlap syndrome (ACOS) has been estimated to be between 13% and 38%^{12,13}. According to new guidelines, this overlap syndrome is characterized by persistent airflow limitation with several features of both asthma and COPD¹⁴. Characteristics of ACOS has not been fully understood due to challenges in defining its clinical features and differences in diagnosis¹⁵.

Many e-cigarette users also use cigarettes, making it difficult to untangle the unique contributions of e-cigarettes to COPD and asthma development. In addition, the long-term respiratory health effects of e-cigarette smoking on ACOS have not been

thoroughly studied, due to challenges of clinical similarities between the two diseases¹⁶. Therefore, we studied the association between e-cigarette use and self-reported asthma, COPD, and ACOS among never combustible cigarette smokers using a large, nationally representative survey in the US to address this gap. We hypothesize that e-cigarette use is associated with chronic respiratory disorders including asthma, COPD and ACOS in adults.

METHODS

Population


We used 2016 (n=486303), 2017 (n=450016), and 2018 (n=437436) data from the Behavioral Risk Factor Surveillance System (BRFSS), a large cross-sectional telephone survey of non-institutionalized adults aged ≥18 years in the US¹⁷. The BRFSS gathers data on health risk behaviors, prevention practices, and access to healthcare as it relates to chronic disorders¹⁸. Our study population consisted of participants, aged ≥18 years, who were never smokers of conventional cigarettes. Participants were considered never smokers of conventional cigarettes if they answered ‘No’ to the question: ‘Have you smoked at least 100 cigarettes in your entire life?’. Therefore, we excluded current or former cigarette smokers (defined as lifetime smoking >100 cigarettes). Participants who did not respond to questions regarding respiratory disorders (asthma, COPD, ACOS) and e-cigarette use were excluded from the analysis (Figure 1). Of 46077 never combustible cigarette smokers, there were 4369 e-cigarette users and 464709 never smokers. After propensity score matching (PSM), the study sample included 4368 e-cigarette users and 4368 never smokers matched on age, sex, race/ethnicity and education level.

Measures

E-cigarette users and never smokers

Respondents who responded ‘Yes’ to the first question ‘Have you ever used an e-cigarette or other electronic vaping product, even just one time, in your entire life?’ were considered as ever cigarette smokers, and those who responded ‘No’ were considered as never e-cigarette smokers. Ever cigarette smokers were then asked: ‘Do you now use e-cigarettes or other electronic vaping products every day, some days, or not at all?’. Those who responded ‘every day’ or ‘some days’ were defined as current e-cigarette users.

Figure 1. Flow chart for participants included in the study



Asthma, COPD, and ACOS

Participants who responded ‘No’ to ‘Has a doctor, nurse, or other health professional ever told you that you have chronic obstructive pulmonary disease or COPD, emphysema, or chronic bronchitis?’ were defined as currently having a diagnosis of asthma if they answered affirmatively to both ‘Has a doctor, nurse, or other health professional ever told you that you have asthma?’ and ‘Do you still have asthma?’. Similarly, respondents who reported to have no asthma were defined as having a diagnosis of COPD if they responded ‘Yes’ to both ‘Has a doctor, nurse, or other health professional ever told you that you have COPD?’ and ‘Do you still have COPD?’. People who responded ‘Yes’ to both asthma and COPD diagnosis were then classified as ACOS.

Sociodemographic and other risk factors

We examined sociodemographic and other risk factors as covariates. The survey included items on age in years (recoded as 5 years intervals from 18 to 80+), gender (female/male), race/ethnicity (non-Hispanic White, non-Hispanic Black, Hispanic, non-Hispanic Multiracial, other), educational level (below high school, graduated from high school, attended college or technical school, graduated from college or technical school), marital status (married, never married or member of unmarried couple, divorced or widowed or separated), income level (<15000, 15000–25000, 25000–35000, 35000–50000, >50000 US\$), employment status (employed, homemaker or student, unemployed), body mass index (BMI) (underweight <18.5, normal weight 18.5–24.9, overweight 25–29.9, and obese >30 kg/m²), and current physical activity or exercise in the last 30 days (Yes/No).

Analysis

The 2016, 2017 and 2018 data were combined and analyzed according to the published Centers for Disease Control and Prevention recommendations, using suggested weighting methodology to improve the representativeness of data¹⁹. Never smokers were matched to e-cigarette smokers on age, sex, race/ethnicity and education using 1:1 PSM to account for selection bias. Specifically, the nearest neighbor matching algorithm was implemented using the MatchIt package²⁰. While there are various methods to employ propensity score matching, nearest neighbor matching is the most common technique that is used in both social sciences and medical literature²¹. This method utilizes a distance measure to quantify the closest match between treated and controlled units. First, propensity scores were estimated based on given covariates using logistic regression. Then, the matching is performed with 1:1 nearest neighbor pair matching without replacement using the difference between the propensity scores of each treated and control unit as a distance measure. For each individual in treatment group (e-cigarette smoker) a control unit (never smoker) was selected using the difference between their propensity score estimates. To assess the balance, standardized difference in mean (SDM) values were calculated. SDMs were smaller than 0.01 and variance ratios were 1 for

all covariates that are used in matching. Potential confounders were identified by creating Directed Acyclic Graphs (DAGs) based on existing literature²². After matching is performed with age, sex and race covariates, the other confounding variables including education level, marital status, income, employment, BMI, and physical activity, were controlled for in the analysis. Multivariable logistic regression was used to examine the association between e-cigarette use and asthma, COPD as well as ACOS. P-values <0.05 were considered significant. All analyses were conducted using R version 3.0.2²³ utilizing survey procedures that account for complex sampling design of BRFSS.


RESULTS

Matched population


The propensity score sample included 4368 e-cigarette smokers matched to 4368 never smokers. Distributional balance before and after PSM for variable age, sex, race/ethnicity and education level, respectively, is shown in Figure 2A. Density plot shows the distributional balance for age, sex, race/ethnicity, and education level before and after the matching. Figure 2B shows covariate balance measured by standardized mean difference indicating standardized difference in means before and after matching on age, sex, and race/ethnicity. For a given variable

Figure 2. Balance diagnostic. A) Distribution balance for matched and unmatched data, B) Covariate balance for variable age, sex, race/ethnicity, and education before and after matching

A. Balance diagnostic for matched and unmatched data



B. Covariate balance for age, sex, race/ethnicity and education level before and after matching



when standardized difference in mean is between the threshold value as indicated by two dashed vertical lines, the balance is considered to be achieved.

Table 1 shows the population weighted baseline distribution of groups after 1:1 propensity score matching on age, sex, race/ethnicity and education


level. In both groups, the age of respondents was 18–24 years. Both groups consisted of a greater proportion non-Hispanic White (64.1%), male (65.6%), and reported to have attended college or technical school (35.9%).

After matching, there were only significant

Table 1. Baseline characteristics of study population after matching on age, sex, race/ethnicity and education level, BRFSS 2016–2018, US (N=8736)

| Characteristics | E-cigarette use status | | p |
|--|------------------------------|------------------------------------|--------|
| | Smokers (N=4368) n (%) | Never smokers (N=4368) n (%) | |
| Age (years) | 18–24 | 18–24 | - |
| Women | 1504 (34.40) | 1504 (34.40) | - |
| Race | | | - |
| White non-Hispanic | 2798 (64.10) | 2798 (64.10) | |
| Black non-Hispanic | 404 (9.20) | 404 (9.20) | |
| Other race non-Hispanic | 345 (7.90) | 345 (7.90) | |
| Multiracial non-Hispanic | 190 (4.30) | 190 (4.30) | |
| Hispanic | 631 (14.40) | 631 (14.40) | |
| Education level | | | - |
| Below high school | 243 (5.60) | 243 (5.60) | |
| Graduated from high school | 1507 (34.50) | 1507 (34.50) | |
| Attended college or technical school | 1567 (35.90) | 1567 (35.90) | |
| Graduated from college or technical school | 1051 (24.10) | 1051 (24.10) | |
| Marital status | | | <0.001 |
| Married | 853 (19.50) | 1354 (30.70) | |
| Never married/cohabiting | 3141 (71.90) | 2760 (63.20) | |
| Divorced/widowed/separated | 374 (8.60) | 267 (6.10) | |
| Income (US\$) | | | 0.726 |
| <15000 | 438 (10.00) | 447 (10.20) | |
| 15000–25000 | 819 (18.80) | 780 (17.90) | |
| 25000–35000 | 518 (11.90) | 498 (11.40) | |
| 35000–50000 | 665 (15.20) | 666 (15.20) | |
| >50000 | 1928 (44.10) | 1977 (45.30) | |
| Employment | | | <0.001 |
| Employed | 2876 (65.80) | 2817 (64.50) | |
| Homemaker/student | 965 (22.10) | 1001 (25.20) | |
| Unemployed | 527 (12.10) | 450 (10.30) | |
| BMI (kg/m²) | | | 0.424 |
| Underweight (<18.5) | 140 (3.20) | 122 (2.80) | |
| Normal weight (18.5–24.9) | 1852 (42.40) | 1832 (41.90) | |
| Overweight (25.0–29.9) | 1327 (30.40) | 1311 (30.00) | |
| Obese (>30.0) | 1049 (24.00) | 1103 (25.30) | |
| Physical activity | | | 0.104 |
| Performs physical activity or exercise | 3690 (84.50) | 3633 (83.20) | |
| No physical activity or exercise in last 30 days | 678 (15.50) | 735 (16.80) | |
| Pulmonary disease | | | <0.001 |
| Asthma | 430 (9.80) | 314 (7.20) | |
| COPD | 90 (2.10) | 58 (1.30) | |
| ACOS | 66 (1.50) | 31 (0.70) | |

Figure 3. Multinomial regression of the association between e-cigarette use and asthma, COPD, and ACOS among adults aged 18–24, BRFSS, 2016, 2018



differences with marital status, employment and pulmonary diseases. A greater proportion of never smokers were married (31.0%) compared with e-cigarette smokers (19.5%). Employment for e-cigarette smokers and never smokers was 65.8% and 66.3%, respectively, with a greater proportion of never smokers reporting an income >50000 US\$ (44.1% vs 46.2%). A greater proportion of e-cigarette smokers reported having pulmonary diseases compared with never smokers. Approximately 9.8% of e-cigarette smokers were diagnosed with asthma compared with 7.2% among never smokers. Also, the proportion of COPD and ACOS was greater among e-cigarette smokers (2.1% vs 1.3%) and (1.5% vs 0.7%), respectively. For all other variables, the differences between the two groups were not significantly different.

Figure 3 shows estimates from multinomial regression analysis after removing participants with missing variables and matching never smokers to e-cigarette smokers on age, sex, race/ethnicity and education level. After adjusting for confounding variables, e-cigarette smokers had increased odds of self-reported ACOS (OR=2.27; 95% CI: 2.23–2.31), asthma (OR=1.26, 95% CI: 1.25–1.27) and COPD (OR=1.44; 95% CI: 1.42–1.46).

DISCUSSION

To our knowledge, this is the first study to examine the association between e-cigarette use and self-reported asthma, COPD, and ACOS among never combustible cigarette smokers using a large nationally representative survey of the US population, i.e. BRFSS. Our findings show that e-cigarette use among adults is associated with chronic respiratory disorders such as asthma, COPD, and ACOS. Using our matched design (i.e. age, sex, race/ethnicity, and education level). The odds of ACOS were twice as high among e-cigarette users compared with never smokers of conventional cigarettes. Data from this large nationally representative sample also showed that e-cigarette users had significantly greater odds of asthma and COPD. The findings from this study suggest the need to further investigate the long-term and short-term health effects since the age of those at risk in our matched design was 18–24 years.

Consistent with other national studies, the majority of adult e-cigarette users were aged 18–24 years^{24,25}. This is not surprising because national studies consistently show higher e-cigarette use among young adults compared to older adults^{26,27}. However, what is unique is that there are adults within these age group reporting COPD and ACOS. Although asthma can

present in younger individuals, including children²⁸, onset of COPD commonly occurs in those over the age of 40 years²⁹. A large international study has recently estimated the prevalence of COPD according to the GOLD standard among young adult population (aged 20–44 years)³⁰. The results from this study support our findings that early onset of COPD-like features likely develop earlier than is usually believed and can inform early detection of the disease and preventive measures. As expected with this young age group, a small percentage of subjects had mild to moderately severe COPD symptoms and none from this age group was found in the most severe stage of COPD (i.e. stage IV). This paves the way for biomarkers such as alpha-1-antitrypsin deficiency to identify younger people who are likely to develop early stages of COPD³¹.

Strengths and limitations

A major strength of our study is the large, representative sample of the US population to test our hypothesis. By selecting never cigarette smokers, we removed the confounding effect attributed to dual use of conventional cigarettes. We also used propensity score matching to remove confounding effects of potential confounders including age, sex, race/ethnicity and education level. Our study is not without limitations. We cannot completely discount residual confounding due to unknown factors. Considering that the BRFSS is a cross-sectional survey design, a causal relationship between e-cigarette use and asthma, COPD, and ACOS cannot be evaluated. Particularly, considering that the age of the study respondents who are e-cigarette users was 18–24 years and that asthma prevalence is higher mostly during childhood³², it is possible that some e-cigarette smokers may have started after asthma diagnosis, ruling out causality. Frequency of self-reported disease outcomes were low in both e-cigarette smokers and matched never smokers. Therefore, it should be noted that the statistically significant difference and more precise odds ratio estimates are observed when the three disease symptoms are combined. We acknowledge that exposure and outcome were self-reported, and there are no data on e-cigarette use initiation, duration, intensity (puffs/day) as well as specific flavor used. Also, we were not able to verify subjects in the two groups using specific biomarkers of exposure. Although we did not imply

a causal association, chronic inhalation of e-cigarette vapor that contains nicotine has been shown to disrupt airway barrier function and induce systemic inflammation in mice³³. Self-reported diagnosis of asthma by a healthcare provider is commonly used by government agencies and scientists in general, as it has been shown to correlate well with the diagnosis of asthma³⁴. However, self-reported physician diagnosis of COPD may have high specificity and low sensitivity³⁵ and hence would bias the effect measure estimate towards the null but this does not explain our finding. Additionally, respiratory disorders are complex disorders resulting from many factors, which may include the interaction between genetic and environmental factors³⁶. In our study, we were not able to account for gene-environment interaction. Despite these limitations, combined data of 2016, 2017, and 2018 from BRFSS, a large nationally representative US sample provided us with a large sample size to study e-cigarette use specifically among this unique population of never combustible cigarette users.

CONCLUSIONS

Data from this large nationally representative sample suggest that e-cigarette use is associated with increased odds of self-reported asthma, COPD, and ACOS among never combustible cigarette smokers. The odds of ACOS were twice as high among e-cigarette users compared with never smokers of conventional cigarettes. The findings from this study suggest the need to further investigate the long-term and short-term health effects of e-cigarette use, since the age of those at risk in our study was 18–24 years. This is not surprising because national studies consistently show higher e-cigarette use among young adults compared to older adults.

REFERENCES

1. Noel JK, Rees VW, Connolly GN. Electronic cigarettes: a new 'tobacco' industry? *Tob Control*. 2011;20(1):81–81. doi:10.1136/tc.2010.038562
2. Grana R, Benowitz N, Glantz SA. E-cigarettes: a scientific review. *Circulation*. 2014;129(19):1972–1986. doi:10.1161/circulationaha.114.007667
3. Goniewicz ML, Smith DM, Edwards KC, et al. Comparison of nicotine and toxicant exposure in users of electronic cigarettes and combustible cigarettes. *JAMA Netw Open*. 2018;1(8):e185937. doi:10.1001/jamanetworkopen.2018.5937
4. Creamer MR, Wang TW, Babb S, et al. Tobacco product

- use and cessation indicators among adults—United States, 2018. *MMWR Morb Mortal Wkly Rep.* 2019;68(45):1013. doi:10.15585/mmwr.mm6845a2
5. National Center for Chronic Disease Prevention and Health Promotion US - Office on Smoking and Health. The health consequences of smoking—50 years of progress: a report of the Surgeon General. Atlanta, GA: Centers for Disease Control and Prevention US; 2014.
 6. Scheffler S, Dieken H, Krischenowski O, Förster C, Branscheid D, Aufderheide M. Evaluation of E-cigarette liquid vapor and mainstream cigarette smoke after direct exposure of primary human bronchial epithelial cells. *Int J Environ Res Public Health.* 2015;12(4):3915-3925. doi:10.3390/ijerph120403915
 7. National Academies of Sciences, Engineering, and Medicine, Health and Medicine Division, Board on Population Health and Public Health Practice, Committee on the Review of the Health Effects of Electronic Nicotine Delivery Systems. Public health consequences of e-cigarettes. Washington, DC: National Academies Press US; 2018. PMID:29894118
 8. Wills TA, Pagano I, Williams RJ, Tam EK. E-cigarette use and respiratory disorder in an adult sample. *Drug Alcohol Depend.* 2019;194:363-370. doi:10.1016/j.drugaledep.2018.10.004
 9. Osei AD, Mirbolouk M, Orimoloye OA, et al. The association between e-cigarette use and asthma among never combustible cigarette smokers: behavioral risk factor surveillance system (BRFSS) 2016 & 2017. *BMC Pulm Med.* 2019;19(1):1-6. doi:10.1186/s12890-019-0950-3
 10. American Lung Association. Asthma Trends and Burden. Accessed November 26, 2020. <https://www.lung.org/research/trends-in-lung-disease/asthma-trends-brief/trends-and-burden>
 11. Ford ES, Croft JB, Mannino DM, Wheaton AG, Zhang X, Giles WH. COPD surveillance—United States, 1999-2011. *Chest.* 2013;144(1):284-305. doi:10.1378/chest.13-0809
 12. Barrecheguren M, Esquinas C, Miravittles M. The asthma-chronic obstructive pulmonary disease overlap syndrome (ACOS): opportunities and challenges. *Curr Opin Pulm Med.* 2015;21(1):74-79. doi:10.1097/MCP.000000000000118
 13. Harada T, Yamasaki A, Fukushima T, et al. Causes of death in patients with asthma and asthma-chronic obstructive pulmonary disease overlap syndrome. *Int J Chron Obstruct Pulmon Dis.* 2015;10:595. doi:10.2147/COPD.S77491
 14. Gibson P, Simpson J. The overlap syndrome of asthma and COPD: what are its features and how important is it? *Thorax.* 2009;64(8):728-735. doi:10.1136/thx.2008.108027
 15. Bateman ED, Reddel HK, van Zyl-Smit RN, Agusti A. The asthma-COPD overlap syndrome: towards a revised taxonomy of chronic airways diseases? *Lancet Respir Med.* 2015;3(9):719-728. doi:10.1016/S2213-2600(15)00254-4
 16. Hines KL, Peebles RS. Management of the asthma-COPD overlap syndrome (ACOS): a review of the evidence. *Curr Allergy Asthma Rep.* 2017;17(3). doi:10.1007/s11882-017-0683-4
 17. Centers for Disease Control and Prevention. Behavioral Risk Factor Surveillance System: 2016 Summary Data Quality Report. https://www.cdc.gov/brfss/annual_data/2016/pdf/2016-sdqr.pdf. Published 2017. Accessed March 2, 2020.
 18. Centers for Disease Control and Prevention. About BRFSS. 2014. Accessed January 25, 2021. <https://www.cdc.gov/brfss/about/index.htm>
 19. Centers for Disease Control Prevention. Behavioral Risk Factor Surveillance System 2016. Weighting BRFSS data: BRFSS 2016. https://www.cdc.gov/brfss/annual_data/2016/pdf/weighting_the_data_webpage_content.pdf. Accessed March 2, 2020.
 20. Ho D, Imai K, King G, Stuart E, Whitworth A. MatchIt: Nonparametric Preprocessing for Parametric Causal Inference [computer program]. Version 3.0.2. 2018.
 21. Austin PC. The performance of different propensity-score methods for estimating differences in proportions (risk differences or absolute risk reductions) in observational studies. *Stat Med.* 2010;29(20):2137-2148. doi:10.1002/sim.3854
 22. Greenland S, Pearl J, Robins JM. Causal diagrams for epidemiologic research. *Epidemiology.* 1999;10(1):37-48. doi:10.1097/00001648-199901000-00008
 23. R Core Team. R: A language and environment for statistical computing. 2013.
 24. Schoenborn CA, Gindi RM. Electronic cigarette use among adults: United States, 2014. *NCHS Data Brief.* 2015;(217). PMID:26555932
 25. Dai H, Leventhal AM. Prevalence of e-cigarette use among adults in the United States, 2014-2018. *JAMA.* 2019;322(18):1824-1827. doi:10.1001/jama.2019.15331
 26. Villarroel MA, Cha AE, Vahratian A. Electronic cigarette use among US adults, 2018. *NCHS Data Brief.* 2020;(365). PMID:32487293
 27. Mayer M, Reyes-Guzman C, Grana R, Choi K, Freedman ND. Demographic Characteristics, Cigarette Smoking, and e-Cigarette Use Among US Adults. *JAMA Netw Open.* 2020;3(10):e2020694. doi:10.1001/jamanetworkopen.2020.20694
 28. Ebmeier S, Thayabaran D, Braithwaite I, Bénamara C, Weatherall M, Beasley R. Trends in international asthma mortality: analysis of data from the WHO Mortality Database from 46 countries (1993-2012). *Lancet.* 2017;390(10098):935-945. doi:10.1016/S0140-6736(17)31448-4
 29. Wheaton AG, Cunningham TJ, Ford ES, Croft JB. Employment and activity limitations among adults with chronic obstructive pulmonary disease—United States, 2013. *MMWR Morb Mortal Wkly Rep.* 2015;64(11):289. PMID:25811677
 30. De Marco R, Accordini S, Cerveri I, et al. An international survey of chronic obstructive pulmonary disease in young adults according to GOLD stages. *Thorax.* 2004;59(2):120-125. doi:10.1136/thorax.2003.011163
 31. Rahaghi FF, Sandhaus RA, Strange C, et al. The prevalence

- of alpha-1 antitrypsin deficiency among patients found to have airflow obstruction. *COPD*. 2012;9(4):352-358. doi:10.3109/15412555.2012.669433
32. Moorman JE, Akinbami LJ, Bailey C, et al. National surveillance of asthma: United States, 2001-2010. *Vital Health Stat 3*. 2012(35):1-58. PMID:24252609
33. Crotty Alexander LE, Drummond CA, Hepokoski M, et al. Chronic inhalation of e-cigarette vapor containing nicotine disrupts airway barrier function and induces systemic inflammation and multiorgan fibrosis in mice. *Am J Physiol Regul Integr Comp Physiol*. 2018;314(6):R834-R847. doi:10.1152/ajpregu.00270.2017
34. Mirabelli MC, Beavers SF, Flanders WD, Chatterjee AB. Reliability in reporting asthma history and age at asthma onset. *J Asthma*. 2014;51(9):956-963. doi:10.3109/02770903.2014.930480
35. Murgia N, Brisman J, Claesson A, Muzi G, Olin AC, Torén K. Validity of a questionnaire-based diagnosis of chronic obstructive pulmonary disease in a general population-based study. *BMC Pulm Med*. 2014;14(1):49. doi:10.1186/1471-2466-14-49
36. Kleeberger SR, Peden D. Gene-environment interactions in asthma and other respiratory diseases. *Annu Rev Med*. 2005;56:383-400. doi:10.1146/annurev.med.56.062904.144908

CONFLICTS OF INTEREST

The authors have completed and submitted the ICMJE Form for Disclosure of Potential Conflicts of Interest and none was reported.

FUNDING

There was no source of funding for this research.

PROVENANCE AND PEER REVIEW

Not commissioned; externally peer reviewed.