



RESEARCH

Open Access

A retrospective analysis of the association between tobacco smoking and deaths from respiratory and cardiovascular diseases in the Kassena-Nankana districts of Northern Ghana

Philip Ayizem Dalinjong^{*}, Paul Welaga, Daniel K Azongo, Samuel Chatio, Dominic Anaseba, Felix Kondayire, James Akazili, Cornelius Debpuur and Abraham Rexford Oduro

Abstract

Background: Tobacco use is a public health problem, responsible for approximately six million deaths annually worldwide. It is a risk factor for many diseases including cancers, respiratory and cardiovascular diseases. In low-and middle-income countries, respiratory and cardiovascular diseases are important causes of death. Tobacco use is prevalent in Ghana, but no study had examined the relationship between tobacco use and deaths from respiratory and cardiovascular diseases in the Upper East Region of Northern Ghana. Hence the paper assessed the association between tobacco use and deaths from respiratory and cardiovascular diseases in that region.

Methods: The study used verbal autopsy data collected from the Kassena-Nankana East and West districts of the Upper East Region of Northern Ghana. Data from deceased individuals aged 15 to 59 years whose deaths occurred between January 1, 2004 and December 31, 2011 and with a known cause as well as smoking status were analyzed. Two binary outcome variables were generated from the cause of death data; whether an individual died from respiratory diseases or not, and from cardiovascular diseases or not. Multiple logistic regression models were used to assess the relationship between tobacco use and deaths from respiratory and cardiovascular diseases.

Results: Out of 3,492 deaths with a known cause of death and smoking status, a third of them smoked. About 16.6% of smokers and 8.1% of non-smokers died from respiratory diseases. Approximately, 10.7% of smokers died from cardiovascular diseases compared to 10.6% of non-smokers. In multivariate analyses, individuals with a history of smoking had two-fold increased odds [OR=2.18, 95% CI (1.6-2.9)] of dying from respiratory diseases. Besides, the number of years of smoking as well as the frequency of smoking is significantly associated with deaths from respiratory diseases. No association existed between tobacco use and deaths from cardiovascular diseases.

Conclusions: Within our study we identified a strong relationship between tobacco use and deaths from respiratory diseases, but not an association with deaths from cardiovascular diseases. Our findings highlight the need to make appropriate health interventions to control tobacco use and thus help reduce premature deaths from respiratory and other tobacco linked diseases.

Keywords: Tobacco, Cigarette smoking, Respiratory diseases, Cardiovascular diseases, Verbal autopsy, Navrongo, Kassena-Nankana districts, Ghana

* Correspondence: padalinjong@yahoo.com
Navrongo Health Research Center, Ghana Health Service, Post Office Box 114,
Navrongo, Ghana, Africa

Introduction

Tobacco use is increasingly a public health problem. Globally, an estimated 6 million people died in 2011 due to smoking [1,2]. It is further projected to kill 50% more people by 2015 than the disease HIV/AIDS [3] and these estimates are expected to rise to 8 million by 2030 if the trend is left unchecked [1]. The burden of tobacco use is gradually shifting from high to low- and middle-income countries [4,5], since the multi-national companies involved in the tobacco industry intend making up for the losses in the advanced countries [2,6,7]. Following this, deaths attributed to tobacco use are expected to reduce by 9% between 2002 and 2030 in the developed countries, but estimated to double from 3.4 million to 6.8 million annually in low- and middle-income countries [3].

Tobacco use is one of the major risk factors for non-communicable diseases [3,8-11] some of which include cardiovascular and respiratory diseases, cancers, etc. Globally, cancer, cardiovascular and chronic respiratory diseases are gauged to be the leading causes of deaths by 2015 [3].

In the context of low- and middle-income countries, cardiovascular diseases are important causes of deaths [12]. According to the WHO, over 80% of the global deaths from cardiovascular diseases occur in low- and middle-income countries, disproportionately affecting the poorest in these settings [13]. A verbal autopsy approach in Vietnam demonstrated that cardiovascular diseases were the leading cause of death in 1999 [14]. Respiratory diseases are also noted to have caused a devastating effect in Africa [15]. Indeed, tobacco users in Africa are expected to die 20 to 25 years earlier from various cancers, respiratory diseases, as well as cardiovascular diseases [16].

Tobacco use is prevalent in Ghana. The common forms of usage include not only cigarette smoking, but also pipe smoking, chewing, sniffing and oral or nasal use [17]. Until recently, there were no studies showing tobacco prevalence for the whole country Ghana [18], most prevalence studies were conducted in the southern urban areas. However, a recent study using data from the 2008 Ghana Demographic and Health Survey found the prevalence of cigarette smoking and other tobacco products to be 7.9% for men and 0.4% for women [18]. As it is in other countries, the evidence demonstrated that tobacco use in Ghana is high among men, rural residents, the poor and those with no formal education [7,18,19].

Geographically, the three regions of Northern Ghana which are considered the poorest, lead in the prevalence rates for cigarette and other tobacco products usage; 31.2% for Upper East Region, 22.5% for Northern Region, and 7.9% for Upper West Region [19]. Despite this high prevalence, no study has documented the relationship between tobacco use and deaths from respiratory and cardiovascular diseases in Northern Ghana, especially

the Upper East Region which had the highest prevalence rate. This paper used verbal autopsy data to explore the relationship between tobacco use and deaths from respiratory and cardiovascular diseases among adults aged 15 to 59 years in the Kassena-Nankana East and West districts of the Upper East Region of Northern Ghana.

Verbal autopsy data is used against the background that certification of deaths in Africa is uncommon. According to Adjuik et al, the registration of deaths in Africa is less than 10% compared to the situation in Europe which is almost 100% [20]. Besides, access to health care services is limited and most deaths take place outside health facilities, and thus very challenging to ascertain the cause of death for each individual. But certification of deaths assist to understand the key causes of death in order to take appropriate policy action or implement interventions directed to specific causes. Verbal autopsy is an innovative way of understanding the causes of deaths in the absence of official certification. It is an interview tool that is used in various places and settings to collect data on causes of death [21], especially in settings with poor vital statistics and weak health systems. This paper contributes to our understanding of the relationship between smoking and dying from respiratory and cardiovascular diseases in the study area and other similar settings. This is essential for the development of effective and efficient health policies on tobacco use and the evaluation of existing programs and policies. It will also provide justification for spending on policies on tobacco use by the Government of Ghana.

Methods

Study design and setting

The study was cross sectional and carried out in the Kassena-Nankana East and West districts (previously the Kassena-Nankana districts) of the Upper East Region of Northern Ghana. There are approximately 153, 293 residents in the two districts [22]. The districts have a total land area of 1,674 square kilometers and mainly covered by the Sahel and Sudan-Savannah types of vegetation. Topographically, the land area is low-lying with an average height of 1,000 meters above sea level. Average rainfall per annum is 950 mm. In all, there are 216 communities in the districts, of which majority are rural, with only 13% of the population living in urban areas. The main economic activity for the people is agriculture (70%). The districts are considered among the poorest in Ghana, due to their reliance on subsistence agriculture.

Existing health facilities include 6 health centres, a district hospital, 27 community-based health and planning services (CHPS) compounds, 2 faith-based health facilities and 3 private clinics. In 2007, the doctor-patient ratio was 1:31927 [23]. Malaria, respiratory infections, skin

and diarrheal diseases were the four leading causes of death for the year 2008 [23].

Data

The study used verbal autopsy data to determine the association between tobacco use and deaths from respiratory and cardiovascular diseases. The verbal autopsy data was collected by the Navrongo Health Research Centre through the Health and Demographic Surveillance System (HDSS). The HDSS is a community registration system that collects and updates longitudinal data on births, deaths, pregnancies, marriages and migrations every four months. The HDSS also conducts verbal autopsy for all deaths registered in the surveillance system [22,24]. In 2003, the verbal autopsy tool was modified to collect additional information on the lifestyle of all deceased individuals including their drinking and smoking patterns. The verbal autopsy instrument has both open-and close-ended questions that document information on the signs, symptoms, lifestyle (smoking and drinking patterns) and circumstances leading to death [25,26]. The tool includes a section for verbatim narrations of the circumstances leading to death. The data is usually collected through interviews with close relatives of the deceased using the verbal autopsy instrument. To minimize recall bias by respondents, the interviews are conducted on average 3 months after a death has occurred. Experienced field supervisors are used to carry out the interviews.

Upon completion of the verbal autopsy questionnaire forms, the next stage involved the assignment of possible cause of death by physicians. Three physicians independently review each verbal autopsy questionnaire form and assign an underlying cause of death corresponding to the 3-digit code of the international statistical classification of diseases and health-related problems [27]. If at least two agree on the underlying cause of death, a diagnosis was established. Where there was disagreement among all three, the form was submitted to two additional physicians for review. Where there was verbal autopsy information but no underlying cause of death could be agreed on, the case was declared undetermined. Where little or no information was available to enable an assignment of cause of death, the diagnosis was declared unknown/undetermined.

Measurement of smoking among the deceased

In measuring the smoking status of the deceased, questions are asked a close relative who provided care for the deceased prior to death, whether the deceased ever used or smoked tobacco during his/her life time. If the response is “yes”, then further questions are asked on the duration of smoking/use of tobacco, frequency, quantity of cigarette smoked/tobacco used, and type of cigarette/tobacco consumed.

Variables

The main exposure variable for the analysis was the smoking status of the deceased prior to death. On the other hand, the outcome variables for the study were respiratory and cardiovascular diseases. First we determined whether an individual died from respiratory disease or not. If an individual died from respiratory disease, it was categorized as 1, and 0 if otherwise. For this analysis, we classified the following diseases under the category of respiratory diseases; chronic obstructive lung disease, tuberculosis, asthma, acute respiratory infections (including pneumonia and influenza), and other respiratory system illness (including empyema, fistula etc.). The second outcome variable was whether an individual died from cardiovascular disease or not, (1, if the death was associated with cardiovascular disease, and 0 if otherwise). The following diseases were also classified as cardiovascular diseases; hypertensive heart disease, ischemic heart disease, cerebrovascular disease and all other cardiovascular diseases. Other diseases not related to either respiratory or cardiovascular diseases were categorized as “other”.

Data analysis

The analysis was done using STATA 11.2 [28]. All deaths aged 15–59 years and occurring between January 1, 2004 and December 31, 2011 were analyzed. Individuals were more likely to initiate smoking at age 15 and thus the reason for using the age group 15 years and above. Besides, the age range 15–59 years was considered the productive age group. Therefore deaths occurring between that age group imply a greater loss of life expectancy, having a substantial impact on society. Overall 4,294 adult deaths occurred in the specified period. Figure 1 shows the distribution of deaths for the period.

Separate multiple logistic regression models were utilized to assess the relationship between tobacco use and dying from respiratory diseases, and again between tobacco use and dying from cardiovascular diseases. Adjusted odds ratios (OR) with 95% confidence intervals (CIs) were calculated. Reference categories were defined as those usually associated with the lowest risk of dying from respiratory or cardiovascular diseases. Potential confounding variables controlled for were age, sex, place of residence and socio-economic status using wealth index generated from household assets using principal component analysis (PCA). The potential confounding variables were arrived at based on their association with tobacco use and dying from respiratory or cardiovascular diseases from previous studies. Besides, socio-economic status was classified as poorest, poorer, poor, less poor, and least poor. Assets used for the socio-economic computation included both large (for example, land and car ownership) and small household items comprising of radio sets, electric fans, etc.




Figure 1 Distribution of deaths for the period 2004–2011.

Ethics approval

Verbal consent was obtained from household heads or any other senior member of the household before carrying out each update for the health and demographic surveillance system. The update for household members was done every four months which includes the collection of the verbal autopsy data. This consent process had been approved by the Institutional Review Board (IRB) of the Navrongo Health Research Centre.

Results

Background characteristics of deaths from respiratory and cardiovascular diseases

Table 1 describes the percentage distribution of the background characteristics of the deaths. Among the deaths with a known smoking status (3,492), approximately 31.7% (1,106) of them were reported to be tobacco smokers preceding their death. About 16.6% of tobacco smokers and 8.1% of non-smokers died from respiratory diseases. Among the smokers, 10.7% died from cardiovascular diseases compared to 10.6% of non-smokers. About 50.8% of men and 2.2% of women were tobacco smokers. Around 12.3% of men and 8.5% of women died from respiratory diseases. There was no marked difference in cardiovascular deaths by sex with 10.4% of men and 11.1% of women

dying from cardiovascular diseases. The median age of the deceased was 45 years (IQR: 35–53).

Association between tobacco use and deaths from respiratory diseases

Table 2 shows the relationship between smoking and dying from respiratory diseases. Individuals who smoked were associated with more than two-fold increased odds of dying from respiratory diseases, in bivariate [OR = 2.31, 95% CI (1.9-2.9)] and multivariate analysis [OR = 2.18, 95% CI (1.6-2.9)]. Likewise, the bivariate analysis indicated that, being older than 30 years, male sex, living in a rural area and a lower socio-economic status was associated with increased odds of dying from respiratory diseases. However, after adjustment for other confounding variables, it was only age that remained significant. Individuals in the age group 50–59 had nearly two-fold increased odds [OR = 1.99, 95% CI (1.3-3.0)] of dying from respiratory diseases. There was no significant association between educational achievement and dying from respiratory diseases.

Association between tobacco use and deaths from cardiovascular diseases

Table 3 demonstrates the association between tobacco use and deaths from cardiovascular diseases. From the Table 3, there was no statistical significance for tobacco use and

Table 1 Percentage distribution of background characteristics of deaths from respiratory and cardiovascular diseases

Variable	Number (%)	Respiratory diseases	Cardiovascular diseases
	N=3,492	n (%)	n (%)
Smoking status			
Smokers	1,106 (31.7)	184 (16.6)	118 (10.7)
Non-smokers	2,386 (68.3)	193 (8.1)	253 (10.6)
Age group			
15-29	560 (16.0)	33 (5.9)	19 (3.4)
30-49	1,604 (45.9)	160 (10.0)	137 (8.5)
50-59	1,328 (38.0)	184 (13.9)	215 (16.2)
Sex			
Female	1,367 (39.2)	116 (8.5)	151 (11.1)
Male	2,125 (60.9)	261 (12.3)	220 (10.4)
Education			
No education	1,584 (45.4)	175 (11.1)	194 (12.3)
Primary/JSS	1,336 (38.3)	150 (11.2)	108 (8.1)
Secondary/tertiary	391 (11.2)	30 (7.7)	51 (13.0)
Missing	180(5.2)	22 (12.2)	18 (10.0)
Place of residence			
Rural	2,970 (85.1)	340 (11.5)	302 (10.2)
Urban	522 (15.90)	37 (7.1)	69 (13.2)
Socio-economic status			
Poorest	751 (21.6)	85 (11.3)	77 (10.2)
Poorer	729 (21.0)	87 (11.9)	71 (9.7)
Poor	749 (21.6)	100 (13.4)	69 (9.2)
Less poor	763 (22.0)	70 (9.2)	83 (10.9)
Least poor	483 (13.9)	4 (7.0)	69 (14.3)
Frequency of smoking			
Chain or hourly smoking	177 (5.1)	37 (20.9)	17 (9.6)
Daily smoking	858 (24.6)	138 (16.1)	90 (10.5)
Occasional smoking	71 (2.0)	9 (12.7)	11 (15.5)
No smoking	2,386 (68.3)	193 (8.1)	253 (10.6)
Duration of smoking			
1-5 years	531 (15.2)	81 (15.3)	62 (11.7)
6-15 years	73 (2.1)	11 (15.1)	8 (11.0)
>15 years	502 (14.4)	92 (18.3)	48 (9.6)
Non-smokers	2,386 (68.3)	193 (8.1)	253 (10.6)
Total	3,492	377 (10.8)	371 (10.6)

deaths from cardiovascular diseases. But in multivariate analysis, being older than 30 years and living in an urban setting was associated with increased odds of dying from cardiovascular diseases. For instance, the age group 50–59 years had over five-fold increased odds [OR = 5.51, 95%

CI (3.1-9.2)] of dying from cardiovascular diseases compared to the age group 15–29. In addition, urban dwellers had more than one-fold increased odds [OR = 1.44, 95% CI (1.1-1.9)] of dying from cardiovascular diseases compared to rural dwellers.

Relationship between length, frequency of smoking and dying from respiratory diseases

Individuals who smoked were further classified into groups based on the frequency of smoking and the duration of smoking. Table 4 describes the relationship between frequency of smoking and dying from respiratory diseases, and between duration of smoking and dying from respiratory diseases. This has been examined in two separate regression models. The odds of dying from respiratory diseases increases based on the frequency of smoking. For example, the multivariate analysis showed that chain/hourly smokers had nearly three-fold increased odds [OR = 2.97, 95% CI (1.9-4.6)] of dying from respiratory diseases compared to non-smokers. Those who smoked daily had double the odds of dying from respiratory diseases compared to non-smokers. Occasional smokers were also more likely to die from respiratory diseases compared to non-smokers, but this was not statistically significant.

In addition, the number of years of smoking was significantly associated with dying from respiratory diseases. The odds of dying from respiratory diseases increases with increasing number of years an individual smoked. In the multivariate analysis, for instance, individuals who smoked for over 15 years had over two-fold increased odds [OR = 2.32, 95% CI (1.6-3.2)] of dying from respiratory diseases compared to non-smokers.

Discussion

This study has illustrated that among the deaths recorded for the period in the study area, 31.7% smoked. This is consistent with the results of a recent study conducted in Ghana. Data from the World Health Organization's (WHO) multi-country study on global aging and adult health (SAGE) was analyzed and it revealed that tobacco prevalence in the Upper East Region (the region for our study), was 31.2% [19]. The finding might be explained by the fact that rural areas usually have high prevalence rates for tobacco use compared to semi-urban and urban places [29,30]. For example, Chockalingam et al. reported that tobacco use was significantly higher for rural residence (23.7%) compared to semi-urban (20.9%) and urban locations (19.4%) [31].

The study also demonstrated that smokers who died from respiratory and cardiovascular diseases were 16.6% and 10.7% respectively. This correlates with other studies as well. Tobacco linked respiratory diseases were shown to be responsible for about 22.1% of deaths in Canada for the year 1998 [32]. In the case of cardiovascular diseases, Pérez

Table 2 Association between smoking and dying from respiratory diseases among deceased adults (n=3,492)

Variable	Unadjusted OR (95% CI)	P-value [#]	*Adjusted OR (95% CI)	P-value [#]
Smoking status				
Smokers	2.31 (1.9-2.9)	<0.001	2.18 (1.6-2.9)	<0.001
Non-smokers	1		1	
Age group				
15-29	1		1	
30-49	1.8 (1.2-2.7)	<0.001	1.35 (0.8-2.0)	0.002
50-59	2.63 (1.8-3.9)		1.99 (1.3-3.0)	
Sex				
Female	1		1	
Male	1.51 (1.2-1.9)	0.006	0.98 (0.7-1.3)	0.904
Education				
No education	1.49 (0.9-2.2)			
Primary/JSS	1.52 (1.0-2.3)	0.117	NS	NS
Secondary/tertiary	1			
Place of residence				
Rural	1.64 (1.1-2.3)	0.007	1.44 (0.9-2.2)	0.081
Urban	1		1	
Socio-economic status				
Poorest	1.61 (1.1-2.5)		1.31 (0.8-2.1)	
Poorer	1.62 (1.1-2.5)	0.009	1.26 (0.8-2.0)	0.216
Poor	2.0 (1.3-3.0)		1.54 (0.9-2.4)	
Less poor	1.29 (0.8-2.0)		1.08 (0.7-1.7)	
Least poor (rich)	1		1	

[#]Overall p-value.

NS – Not significant in bivariate analysis.

*Adjusted for age, sex, place of residence and wealth index in separate models. Missing categories were not included in the regression models.

et al. reported that deaths from cardiovascular diseases among smokers in 1995 and 2007 respectively, were 52% and 42% in Cuba [33]. However, the proportions reported in their study are far higher than that reported in this paper. According to the authors, their finding put Cuba in the lead globally for deaths attributable to tobacco use.

Besides deaths from respiratory diseases were found to be significantly related to smoking as reported in other studies [15,34,35]. In the multivariate analysis, smoking was still found to be a significant risk factor for deaths due to respiratory diseases. This is supported by other studies as well. In Bangladesh, for instance, tobacco use was associated with 1.58 increased odds (99% CI 0.9-2.8) of dying from respiratory diseases [36]. Our findings showed that chained or frequent smokers (people who smoked every hour) had higher chances of dying from respiratory diseases, compared to non-smokers. The results clearly showed a significant dose–response relationship between dying from respiratory diseases and the quantity/

amount of tobacco smoked [37,38]. A similar dose response relationship was observed between the number of years an individual smoked and deaths from respiratory diseases. Thus, the study has demonstrated that tobacco use is a major risk factor for deaths due to respiratory diseases. Therefore tobacco control should be considered a top priority in Ghana.

However, the study could not establish any significant relationship between tobacco use and deaths from cardiovascular diseases. Other studies exist to support our finding. For instance, a prospective study conducted in Bangladesh could not establish any clear association between smoking and deaths from overall cardiovascular diseases [38]. Again, the Singapore Chinese Health Study among men and women of Chinese ethnicity found no apparent relationship between smoking and deaths from stroke [39]. Contrary to our finding, other studies have found a significant association between smoking and deaths from cardiovascular diseases [11,40,41]. Further research

Table 3 Association between smoking and dying from cardiovascular diseases among deceased adults (n = 3,492)

Variable	Unadjusted OR (95% CI)	P-value [#]	*Adjusted OR (95% CI)	P-value [#]
Smoking status				
Smokers	1		1	
Non-smokers	1.02 (0.8-1.3)	0.817	1.16 (0.9-1.5)	0.246
Age group				
15-29	1		1	
30-49	2.58 (1.6-4.3)	<0.001	2.59 (1.7-4.9)	<0.001
50-59	5.45 (3.3-8.9)		5.51 (3.1-9.2)	
Sex				
Female	1.11 (0.9-1.4)	0.357	NS	NS
Male	1			
Education				
No education	1.58 (1.2-2.0)		1.21 (0.9-1.6)	
Primary/JSS	1	0.004	1	0.080
Secondary/tertiary	1.70 (1.2-2.4)		1.49 (1.0-2.2)	
Place of residence				
Rural	1		1	
Urban	1.32 (0.9-1.8)	0.058	1.44 (1.1-1.9)	0.020
Socio-economic status				
Poorest	1.09 (0.8-1.6)			
Poorer	1.08 (0.8-1.5)			
Poor	1	0.151	NS	NS
Less poor	1.22 (0.9-1.7)			
Least poor (rich)	1.56 (1.1-2.3)			

[#]Overall p-value.

NS – Not significant in bivariate analysis.

*Adjusted for age, sex, place of residence and wealth index in separate models.

Table 4 Relative odds of dying from respiratory diseases by frequency and duration of smoking

Variable	Unadjusted OR (95% CI)	P-value [#]	Adjusted* OR (95% CI)	P-value [#]
Frequency of smoking				
Chain/hourly smoking	3.19 (2.2-4.7)		2.97 (1.9-4.6)	
Daily smoking	2.18 (1.7-2.8)	<0.001	2.06 (1.5-2.8)	<0.001
Occasional smoking	1.79 (0.9-3.7)		1.67 (0.8-3.5)	
Non-smokers	1		1	
Duration of smoking				
1-5 years	2.13 (1.6-2.8)		2.13 (1.5-2.9)	
6-15 years	2.12 (1.1-4.1)	<0.001	1.87 (0.9-3.7)	<0.001
>15 years	2.54 (1.9-3.4)		2.32 (1.6-3.2)	
Non-smokers	1		1	

[#]Overall p-value; *Adjusted for age, sex, place of residence and wealth index in separate models.

in that direction is recommended, especially from other INDEPTH Network sites which collect verbal autopsy data.

Age especially is one of the most important factors determining deaths from cardiovascular diseases. From this study, older individuals as well as individuals living in urban areas had increased odds of dying from cardiovascular diseases. Similar findings exist elsewhere to support our finding. Jiang et al. found that deaths resulting from coronary heart disease in Tianjin (China) were higher among older age groups and individuals living in urban areas [42]. However, it is worrying to realize that people in their prime ages or older are dying due to cardiovascular diseases. The impact of these deaths are enormous on the family, employer and the community due to the fact that these people are either the bread winners for their families, the most experienced employees, or sometimes leaders of their communities [43]. There is the need for education on lifestyle habits (for example, dieting, alcohol consumption, physical inactivity, etc.) especially for older individuals as well as urban dwellers.

Study limitations

The use of the verbal autopsy tool for data collection has its limitations including accuracy and recall bias by the respondents. It is a challenge determining whether the caregiver responding for the deceased is truthful in his/her responses. Again, it is possible for non-smokers to have died from tobacco related diseases (both respiratory and cardiovascular) due to exposure to second hand smoke. Our study would not have captured such deaths since the main exposure variable for the analysis was the smoking status of the individual prior to his/her death. Therefore our findings on the association between smoking and deaths from both respiratory and cardiovascular diseases could have been underreported.

Nonetheless, the study was a population based study with vigorous study oversight, well trained field staff for data collection, and a well-defined health and demographic surveillance system for monitoring deaths and other key events. Again, the verbal autopsy tool for the study was developed and standardized by the WHO and the International Network for the Demographic Evaluation of Populations and Their Health (INDEPTH Network).

Policy recommendations

In most sub-Saharan African countries including Ghana, the challenges of tobacco use are emerging, but stringent policies on smoking and use of tobacco products are minimal and not effectively enforced. From the findings of this study, it is clear that smoking is a significant predictor of deaths from respiratory diseases. This therefore, calls for urgent need to re-assess the state of the implementation of the Framework Convention on Tobacco Control

(FCTC) to reduce tobacco use among the adult population. In Ghana particularly, the Public Health Bill incorporating the Tobacco Control Bill requires quick action to become a Legislative Instrument (LI) to allow for its enforceability. The provisions of the Tobacco Control Bill if well enforced will contribute to reducing respiratory and other tobacco related diseases, thus minimizing premature deaths in the country. Following our findings from this study, which took place in a predominantly rural area, there is the need to enact policies addressing tobacco use in both rural and urban areas.

Conclusions

The analysis showed an association between tobacco use and deaths from respiratory diseases in the Kassena-Nankana East and West districts. A significant relationship was observed for tobacco use and deaths resulting from respiratory diseases. Tobacco users were more prone to dying from respiratory diseases compared to non-users. Our study could not establish any significant relationship between tobacco use and deaths from cardiovascular diseases. Only age and urban location were found to be strong predictors of deaths from cardiovascular diseases. There is need to adopt appropriate health interventions to control tobacco use. Controlling tobacco use can contribute to reducing respiratory and other tobacco linked diseases and by extension deaths related to tobacco usage.

Abbreviations

CI: Confidence Interval; FCTC: Framework Convention on Tobacco Control; HDSS: Health and Demographic Surveillance System; LI: legislative instrument; INDEPTH Network: International Network for the Demographic Evaluation of Populations and Their Health; IRB: Institutional Review Board; OR: Odds Ratio; PCA: Principal Component Analysis; WHO: World Health Organization.

Competing interests

The authors declare that they have no competing interests.

Authors' contributions

PAD and PW contributed in the conception, design, analysis, interpretation and drafting of the manuscript. DKA, SC, DA, and FK contributed in the acquisition of data, analysis, interpretation and drafting of the manuscript. JA, CD and ARO contributed in the drafting of the manuscript. All authors read and approved the final copy of the manuscript.

Acknowledgments

The Health and Demographic Surveillance System under the Navrongo Health Research Centre collected the data for this analysis. Therefore, we thank both the field and Computer Centre staff for all their efforts in collecting and managing the data. Funding for the data analysis was provided by the School of Economics, University of Cape Town, South Africa. Professor Corne Van Walbeek and Dr. Evan Blecher provided mentorship for the project. We are very grateful.

Received: 20 September 2014 Accepted: 17 April 2015

Published online: 26 April 2015

References

1. Eriksen M, Mackay J, Ross H. The tobacco atlas. 4th edition. Atlanta: GA: American Cancer Society. New York: NY: World Lung Foundation; 2012.

2. WHO. WHO report on the global tobacco epidemic 2013, enforcing bans on tobacco advertising, promotion and sponsorship. Geneva. Geneva, Switzerland: World Health Organization; 2013.
3. Mathers CD, Loncar D. Projections of global mortality and burden of disease from 2002 to 2030. *PLoS Med*. 2006;3:e442.
4. Jha P, Ranson M, Nguyen S, Yach D. Estimates of global and regional smoking prevalence in 1995, by age and sex. *Am J Public Health*. 2002;92:1002–6.
5. Norberg M, Lundqvist G, Nilsson M, Gilljam H, Weinehall L. Changing patterns of tobacco use in a middle-aged population: the role of snus, gender, age, and education. *Glob Health Action*. 2011;4:10.3402/gha.v4i0.5613.
6. Baleta A. Africa's struggle to be smoke free. *Lancet*. 2010;375:107–8.
7. Doku D, Darteh EK, Kumi-Kyereme A. Socioeconomic inequalities in cigarette smoking among men: evidence from the 2003 and 2008 Ghana Demographic and Health surveys. *Arch Public Health*. 2013;71:9.
8. Thakur J, Garg R, Narain J, Menabde N. Tobacco use: a major risk factor for non-communicable diseases in South-East Asia Region. *Indian J Public Health*. 2011;55:155–60.
9. Kim HC, Oh SM. Noncommunicable diseases: current status of major modifiable risk factors in Korea. *J Prev Med Public Health*. 2013;46:165–72.
10. WHO. World Health Organisation fact sheet, updated March 2013, Non-communicable diseases. Geneva: World Health Organization; 2013. http://www.who.int/features/factfiles/noncommunicable_diseases/en/ (accessed on 20 July 2014).
11. USDHHS. The health consequences of smoking—50 years of progress, Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health 2014. Atlanta, GA: A Report of the Surgeon General; 2014.
12. Leeder SRS, Greenberg H, Liu H, Esson K. A race against time: the challenge of cardiovascular disease in developing economies. New York: Columbia University; 2004.
13. WHO. Cardiovascular diseases (CVDs) Fact sheet N°317 Updated March 2013: World Health Organization; 2013. http://www.who.int/features/factfiles/noncommunicable_diseases/en/ (accessed on 20 July 2014)
14. Huong DL, Minh HV, Byass P. Applying verbal autopsy to determine cause of death in rural Vietnam. *Scand J Public Health Suppl*. 2003;62:19–25.
15. Bates M, Mudenda V, Mwaba P, Zumla A. Deaths due to respiratory tract infections in Africa: a review of autopsy studies. *Curr Opin Pulm Med*. 2013;19:229–37.
16. CVJAFRICA. Cardiovascular Topics. *Cardiovasc J Africa* 2013;24 (2).
17. Drope J: *Tobacco Control in Africa: People, Politics and Policies*. Anthem Press; 2011
18. John RM, Mamudu HM, Liber AC. Socioeconomic implications of tobacco use in Ghana. *Nicotine Tob Res*. 2012;14:1205–12.
19. Yawson AE, Baddoo A, Hagan-Seneadza NA, Calys-Tagoe B, Hewlett S, Dako-Gyeke P, et al. Tobacco use in older adults in Ghana: Socio-demographic characteristics, health risks and subjective wellbeing. *BMC Public Health*. 2013;13:979.
20. Adjuik M, Smith T, Clark S, Todd J, Garrib A, Kinfu Y, et al. Cause-specific mortality rates in sub-Saharan Africa and Bangladesh. *Bull World Health Organ*. 2006;84:181–8.
21. Sethi R, General AR. Sample Registration System in India. Presentation at the Regional Consultation on Mortality Statistics, Report of the Consultation. New Delhi: WHO-SEARO; 2007. p. 24–7.
22. Oduro AR, Wak G, Azongo D, Debpuur C, Wontuo P, Kondayire F, et al. Profile of the Navrongo Health and Demographic Surveillance System. *Int J Epidemiol*. 2012;41:968–76.
23. UNDP, Kassena-Nankana District Human Development Report 2010. Resource Endowment, Investment Opportunities and the Attainment of MDGs Accra: United Nations and Development Programme. 2012.
24. Binka FN, Ngom P, Phillips JF, Adazu K, MacLeod BB. Assessing population dynamics in a rural African society: the Navrongo Demographic Surveillance System. *J Biosoc Sci*. 1999;31:375–91.
25. Fauveau V. Assessing probable causes of death without death registration or certificates: a new science? *Bull World Health Organ*. 2006;84:246–7.
26. Setel PW, Whiting DR, Hemed Y, Chandramohan D, Wolfson LJ, Alberti KG, et al. Validity of verbal autopsy procedures for determining cause of death in Tanzania. *Trop Med Int Health*. 2006;11:681–96.
27. WHO, World Health Organization International Statistical Classification of Diseases and Related Health Problems, 10th Revision (ICD 10): World Health Organization; 2010
28. StataCorp. Statistical software, v. 11.2. College station, Texas: StataCorp LP; 2010.
29. Berraho M, Serhier Z, Tachfouti N, Elfakir S, El Rhazi K, Slama K, et al. Burden of smoking in Moroccan rural areas. *East Mediterr Health J*. 2010;16:677–83.
30. Li Q, Hsia J, Yang G. Prevalence of smoking in China in 2010. *N Engl J Med*. 2011;364:2469–70.
31. Chockalingam K, Vedhachalam C, Rangasamy S, Sekar G, Adinarayanan S, Swaminathan S, et al. Prevalence of tobacco use in urban, semi urban and rural areas in and around Chennai city, India. *PLoS One*. 2013;8:e76005.
32. Makomaski Illing EM, Kaiserman MJ. Mortality attributable to tobacco use in Canada and its regions, 1998. *Can J Public Health*. 2004;95:38–44.
33. Pérez PV, Travieso DH, Roche RGG, Gorbea MB, Pérez TR, Fernández SJV. Smoking-attributable mortality in Cuba. *MEDICC review*. 2009;11:3.
34. Kanungo S, Tsuzuki A, Deen J, Lopez A, Rajendran K, Manna B, et al. Use of verbal autopsy to determine mortality patterns in an urban slum in Kolkata, India. *Bull World Health Organ*. 2010;88:667–74.
35. Dogar O, Bullen C, Novotny TE, Siddiqi K. Smoking cessation and respiratory disease in low-income and middle-income countries. *Lancet Respir Med*. 2013;1:e23–4.
36. Alam DS, Jha P, Ramasundarahettige C, Streatfield PK, Niessen LW, Chowdhury MA. Smoking-attributable mortality in Bangladesh: Proportional mortality study. *Bull World Health Organ*. 2013;91:757–64.
37. Juren AJ, Frohlich J, Ignaszewski A. Commonplace to condemned: the discovery that tobacco kills, and how Richard Doll shaped modern smoking cessation practices. *BCM J*. 2012;54:183.
38. Wu F, Chen Y, Parvez F, Segers S, Argos M, Islam T, et al. A prospective study of tobacco smoking and mortality in Bangladesh. *PLoS one*. 2013;8, e58516.
39. Shankar A, Yuan JM, Koh WP, Lee HP, Yu MC. Morbidity and mortality in relation to smoking among women and men of Chinese ethnicity: the Singapore Chinese health study. *Eur J Cancer*. 2008;44:100–9.
40. Groenewald P, Vos T, Norman R, Laubscher R, van Walbeek C, Saloojee Y, et al. South African comparative risk assessment collaborating Group: Estimating the burden of disease attributable to smoking in South Africa in 2000. *S Afr Med J*. 2007;97:674–81.
41. Mallaina P, Lionis C, Rol H, Imperiali R, Burgess A, Nixon M, et al. Smoking cessation and the risk of cardiovascular disease outcomes predicted from established risk scores: results of the cardiovascular risk assessment among smokers in primary care in Europe (CV-ASPIRE) study. *BMC Public Health*. 2013;13:362.
42. Jiang G, Wang D, Li W, Pan Y, Zheng W, Zhang H, et al. Coronary heart disease mortality in China: age, gender, and urban–rural gaps during epidemiological transition. *Revista panamericana de salud pública*. *Pan Am J Public Health*. 2012;31:317–24.
43. Mbewu A, Mbanya JC. Cardiovascular Disease. In *Disease and Mortality in sub-Saharan Africa*. 2nd edition. Edited by Jamison DT, Feachem RG, Makgoba MW, Bos ER, Baingana FK, Hofman KJ, Rogo KO: Washington (DC); 2006.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

