

1 0.37785714

0.00071429

1 0.37785714

1

Year

Source

Brand

Year

0.37785714

DF Type III SS Mean Square F Value Pr > F

0.00071429


0.37785714

33.25 0.0001

0.06 0.8067

33.25 0.0001


Supplemental Figure 1. Average percentage of alkaloids in the tobacco of six filtered and three non-filtered cigarette brands, 1960-1990



Supplemental Figure 2. Average percentage of reconstituted tobacco in the blend of six filtered and three non-filtered cigarette brands, 1960-1990

Source	DF	Sum of Squares	Mean Square	F Value	Pr > F
Model	2	124.0604762	62.0302381	33.40	<.0001
Error	9	16.7161905	1.8573545		
Corrected Total	11	140.7766667			

	R-Square		re	e Coeff Var Root MSE		percent Mean			
	0.	8812	57	8.352490	1.362848		16.316	67	
Sou	rce	DF		Type I SS	Mean Squa	re	F Value	Р	r > F
Bran	ıd	1	36	.05333333	36.053333	33	19.41	0.0	0017
Year	-	1	88	.00714286	88.007142	86	47.38	<.	0001
Sou	rce	DF	Т	ype III SS	Mean Squa	re	F Value	P	r > F
Bran	d	1	36	.05333333	36.053333	33	19.41	0.	0017
Year	-	1	88	.00714286	88.007142	86	47.38	<.	0001




Supplemental Figure 3. Average percentage of expanded tobacco used in the blend of tobacco of popular filtered and non-filtered cigarettes, 1975-1990

Source	DF	Sum of Squares	Mean Square	F Value	Pr > F
Model	2	50.67000000	25.33500000	8.12	0.0269
Error	5	15.60500000	3.12100000		
Corrected Total	7	66.27500000			

R-Square	Coeff Var	Root MSE	percent Mean
0.764542	16.02390	1.766635	11.02500

Brand 1 0.04500000 0.04500000 0.01 0.9091 Year 1 50.62500000 50.62500000 16.22 0.0100	Source	DF	Type I SS	Mean Square	F Value	Pr > F
Year 1 50.62500000 50.62500000 16.22 0.0100	Brand	1	0.04500000	0.04500000	0.01	0.9091
	Year	1	50.62500000	50.62500000	16.22	0.0100

Source	DF	Type III SS	Mean Square	F Value	Pr > F
Brand	1	0.04500000	0.04500000	0.01	0.9091
Year	1	50.62500000	50.62500000	16.22	0.0100




Supplemental Figure 4. Average TPM (mg/cigt) measurements from the smoke of popular filtered and non-filtered cigarettes, 1960-1990

Source	DF	Sum of Squares	Mean Square	F Value	Pr > F
Model	2	196.1058929	98.0529464	18.09	0.0003
Error	11	59.6226786	5.4202435		
Corrected Total	13	255.7285714			

R-Square	Coeff Var	Root MSE	mgtpm Mean
0.766852	11.07133	2.328142	21.02857

Source	DF	Type I SS	Mean Square	F Value	Pr > F
Brand	1	105.3257143	105.3257143	19.43	0.0010
Year	1	90.7801786	90.7801786	16.75	0.0018

Source	DF	Type III SS	Mean Square	F Value	Pr > F
Brand	1	105.3257143	105.3257143	19.43	0.0010
Year	1	90.7801786	90.7801786	16.75	0.0018



Supplemental Figure 5. Average nicotine (mg/cigt) yield//s in the smoke of popular filtered and non-filtered cigarettes, 1960-1990

Source	DF	Sum of Squares	Mean Square	F Value	Pr > F
Model	2	0.24928571	0.12464286	15.61	0.0006
Error	11	0.08785714	0.00798701		
Corrected Total	13	0.33714286			

R-Square	Coeff Var	Root MSE	mgnic Mean	
0.739407	6.799898	0.089370	1.314286	

Source	DF	Type I SS	Mean Square	F Value	Pr > F		
Brand	1	0.23142857	0.23142857	28.98	0.0002		
Year	1	0.01785714	0.01785714	2.24	0.1630		
Source	DF	Type III SS	Mean Square	F Value	Pr > F		
Source Brand		Type III SS 0.23142857	Mean Square 0.23142857	F Value 28.98	Pr > F 0.0002		


0.01785714

2.24 0.1630

0.01785714

1

Year




Supplemental Figure 6. Average rod length (mm) of popular filtered and non-filtered cigarette brands, 1960-1990

Source	DF	Sum of Squares	Mean Square	F Value	Pr > F
Model	2	106.9551786	53.4775893	74.38	<.0001
Error	11	7.9083929	0.7189448		
Corrected Total	13	114.8635714			

R-Square	Coeff Var	Root MSE	mmrod Mean
0.931150	1.264049	0.847906	67.07857

Source	DF	Type I SS	Mean Square	F Value	Pr > F
Brand	1	95.16071429	95.16071429	132.36	<.0001
Year	1	11.79446429	11.79446429	16.41	0.0019
Source	DF	Type III SS	Mean Square	F Value	Pr > F
Source Brand	DF 1	Type III SS 95.16071429	Mean Square 95.16071429	F Value 132.36	Pr > F <.0001



Supplemental Figure 7. Average rod and filter length (mm) of popular filtered cigarettes, 1960-1990

© 2023 Cummings K.M. et al.