




Supplementary file

© 2026 Kim M. et al.

DOI:

10.18332/tid/236889

The content has been provided by the author(s) and has not been reviewed, verified, or endorsed by European Publishing. It may not have undergone peer review. The views, opinions, and recommendations expressed are solely those of the author(s) and do not necessarily reflect the position of European Publishing. European Publishing accepts no responsibility or liability for any consequences arising from the use of, or reliance on, this content.



Supplementary Table 1. Association between smoking cessation duration and prediabetes by non-smoking-related subgroups

Subgroup	Category	Men aOR (95% CI)	Women aOR (95% CI)
Age	30–39	1.038 (1.008–1.069)*	1.003 (0.953–1.057)
	40–49	0.998 (0.981–1.015)	1.045 (1.009–1.083)*
	50–59	0.989 (0.977–1.002)	0.991 (0.960–1.022)
	60–69	0.988 (0.978–0.998)*	1.011 (0.978–1.045)
	70≤	0.997 (0.989–1.005)	0.993 (0.960–1.026)
Body mass index (kg/m ²)	0–22	0.995 (0.986–1.004)	1.014 (0.990–1.038)
	23–24	0.995 (0.985–1.004)	0.997 (0.967–1.029)
	25≤	0.992 (0.983–1.001)	0.998 (0.973–1.023)
Income group	Low	0.996 (0.985–1.006)	1.009 (0.979–1.039)
	Mid-Low	0.996 (0.986–1.007)	1.009 (0.978–1.041)
	Mid-High	0.997 (0.985–1.008)	1.016 (0.986–1.047)
	High	0.990 (0.979–1.000)*	0.994 (0.958–1.032)
Employment status	Employed	0.994 (0.987–1.000)	1.000 (0.979–1.022)
	Unemployed	0.996 (0.987–1.004)	1.006 (0.985–1.028)
Marital status	Ever married	0.993 (0.988–0.999)*	1.004 (0.989–1.019)
	Never married	1.008 (0.977–1.040)	0.990 (0.908–1.079)
Alcohol consumption	0–4 drinks/sitting	0.993 (0.987–1.000)*	1.008 (0.991–1.025)
	≥5 drinks/sitting	0.994 (0.985–1.003)	1.000 (0.969–1.033)
Aerobic activity	No	0.990 (0.983–0.997)*	1.000 (0.982–1.018)
	Yes	1.000 (0.991–1.008)	1.015 (0.988–1.042)
Hypertension	No	0.993 (0.986–1.000)*	1.010 (0.993–1.028)
	Yes	0.995 (0.987–1.004)	0.994 (0.965–1.024)

aOR, adjusted odds ratio; CI, confidence interval.

* $p < 0.05$